

THE “TWIN JET” PLANETARY NEBULA M2–9

Mario Livio¹ and Noam Soker²

ABSTRACT

We present a model for the structure, temporal behavior, and evolutionary status of the bipolar nebula M2–9. According to this model the system consists of an AGB or post-AGB star and a hot white dwarf companion, with an orbital period of about 120 years. The white dwarf has undergone a symbiotic nova eruption about 1200 years ago, followed by a supersoft x-ray source phase. The positional shift of the bright knots in the inner nebular lobes is explained in terms of a revolving ionizing source. We show that the interaction between the slow, AGB star’s wind, and a collimated fast wind from the white dwarf clears a path for the ionizing radiation in one direction, while the radiation is attenuated in others. This results in the mirror-symmetric (as opposed to the more common point-symmetric) shift in the knots. We show that M2–9 provides an important evolutionary link among planetary nebulae with binary central stars, symbiotic systems, and supersoft x-ray sources.

Subject headings: planetary nebulae: general – planetary nebulae: individual: M2–9 – stars: binaries: close – stars: AGB and post-AGB – stars: mass loss – stars: symbiotic – X-rays: stars

¹Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; mlivio@stsci.edu

²Department of Physics, University of Haifa at Oranim, Oranim, Tivon 36006, Israel; soker@physics.technion.ac.il

1. INTRODUCTION

M2–9 is a spectacular, well studied bipolar nebula, usually classified as a planetary nebula (PN). It consists of two oppositely directed, narrow, cylindrical-like bright lobes, each extending to an angular projected distance of $\sim 20''$ from the center of the nebula (Fig. 1a). Farther out, there are two “ansae,” one on each side along the symmetry axis (Schwarz et al. 1997), at $\sim 60''$ from the center, moving outward at $\sim 190 \text{ km s}^{-1}$ (Schwarz et al. 1997). The inner lobes on the other hand, expand at a much slower speed, of $\sim 40 \text{ km s}^{-1}$ (Schwarz et al. 1997; Solf 2000), or maybe even 15 km s^{-1} (Trammell, Goodrich & Dinerstein 1995; hereafter TGD). The ansae have a point symmetrical structure (namely, are at symmetrical points when reflected about the central star).

The nebula and its central region show many interesting properties, which are summarized in, e.g. Balick (1989), TGD, Hora & Latter (1994), Phillips & Cuesta (1999), Solf (2000), and Doyle et al. (2000; hereafter DBCS). Balick (1989), for example, finds an $\text{H}\alpha$ linewidth of $11,000 \text{ km s}^{-1}$, implying the presence of a wind having a velocity of $\sim 5,500 \text{ km s}^{-1}$ from the nucleus. Balick further notes that “the spectrum of M2–9’s nucleus is more similar to the slow nova RR Tel, some symbiotic stars, and Seyfert (type 1.9) galactic nuclei...” (see also Solf 2000).

In view of these, and other intriguing properties (see below), it is important to identify the nature of this nebula and to place it in a broader evolutionary context. In the present paper we attempt to do precisely that. We also address specifically a perplexing positional (time dependent) shift of the bright knots in the inner lobes, $\sim 8''$ from the nucleus (Allen & Swings 1972), which was recently studied in detail by DBCS. The shifts in the northern and southern lobes are in the *same* direction, i.e. there is a mirror-symmetry rather than a point-symmetry of the bright knots. The picture that emerges from many studies (e.g. Allen & Swings 1972; van den Bergh 1974; Goodrich 1991; Kohoutek & Surdej 1980; and for more information see §1 of TGD) is that of two UV illuminating beams in a mirror symmetry, which move around the central UV source at a period of ~ 100 yrs. TGD conducted a detailed study of the ionization state of the shifting knots. One of their conclusions was that the off-knots regions are ionized by attenuated UV radiation. They assumed that the lobes contain rings of higher density, and concluded that: “The UV source illuminates a section of the lobe, causing the rings of material to glow, producing the knots and explaining and naturally leading to their “fixed” north-south positions. Obscuring material near the star keeps one side of the lobe from being illuminated by the full force of UV source.” TGD also estimated that the column density of the obscuring material is $\sim 10^{18-18.4} \text{ cm}^{-2}$.

Goodrich (1991; see also Scarrott, Scarrott & Wolstencroft 1993) suggested that the radiation that ionizes the knots comes from a companion, and not from the central star itself. DBCS, on the other hand, attribute the ionization to a fast jet hitting the bright spots. As we shall explain in the Discussion (§4), we identify some difficulties in the scenario proposed by DBCS. Instead, we propose in the present paper a different model, which characterizes M2–9 as a stage in the evolution of interacting, detached binary systems.

The important properties of M2–9, as they can be deduced from the different studies are as follows.

1. The bolometric luminosity of the central star is not known. The star is heavily obscured (e.g. no point source is observed in the snapshot HST survey; H. Bond 2000, private communication). A fit to the IR spectrum gives a total flux of $1.1 \times 10^4 L_{\odot} (D/\text{kpc})^2$ (S. Kwok 2000, private communication), but the distance is rather uncertain (Schwarz et al. 1997 suggest ~ 650 pc).
2. Lim & Kwok (2000) argue that the mass loss rate from the central source is $\ll 10^{-5} M_{\odot} \text{ yr}^{-1}$, and maybe even below $10^{-6} M_{\odot} \text{ yr}^{-1}$.
3. The two ansae at very large distances from the lobes expand at $\sim 200 \text{ km s}^{-1}$ (Schwarz et al. 1997), a high velocity compared with that of the lobes which expand at $15\text{--}40 \text{ km s}^{-1}$ (Solf 2000).
4. The ionizing source is very probably a WD companion (e.g. Goodrich 1991; Schwarz et al. 1997; Lim & Kwok 2000).
5. Material moving at $\gtrsim 5,000 \text{ km s}^{-1}$ is found near the center of the nebula (Balick 1989). This wind was suggested to result from an accretion disk around the WD companion (e.g. Torres-Peimbert & Arrieta 1998). This suggestion is consistent with the general finding that outflow speeds are always of the order of the Keplerian speed around the central object (Livio 1999, 2000).
6. The ionizing radiation escapes mainly in two directions, north and south, with an inclination of $\sim 30^{\circ}$ to the symmetry axis. The inclination to the symmetry axis is in the *same* direction for both “beams” (west for both, or east for both; Fig. 1a). The beams rotate around the central star with a period of 120 yrs (DBCS).
7. The ionizing radiation does reach other regions around the lobes, but it is attenuated in those directions (TGD).
8. The Galactic latitude of M2–9 is the largest in the list of 43 bipolar PNe of Corradi & Schwarz (1995). For an assumed distance of 650 pc it implies a Galactic height of 200 pc. This may imply that the progenitor of M2–9 was not very massive compared to other bipolar PNe progenitors (perhaps a late A or F rather than a B star).

We argue that *all* of these properties can be adequately explained in the context of a model in which a central binary system that is a supersoft X-ray source (SXS), is in the transition phase from a symbiotic nebula (possibly generated by a symbiotic nova; Solf 2000) to a PN. In particular, we demonstrate that a system consisting of an asymptotic giant branch (AGB) or a post-AGB star (with its associated slow wind) and a hot WD (which has a fast wind), in an orbital period of

~ 120 yrs, can ionize the “walls” of the lobes in one direction, while the ionization is substantially attenuated in other directions. The revolving ionizing source is explained by a combination of the attenuating effects of the post-AGB star’s wind and by the effects of a collimated fast wind (CFW) blown by the WD companion, which is bent by the slower wind from the post-AGB star. This type of flow, where the CFW is being strongly pushed upon and bent by the AGB (or post-AGB) wind is termed a ‘weak-CFW’ flow by Soker & Rappaport (2000, hereafter SR00). The bent “jets” can both compress the ring in each lobe, and clear a way for the ionizing radiation in the two specific directions around the symmetry axis. The circumbinary material has special properties (in terms of its optical depth, see below) for a relatively short period of time (of the order of a few hundred years), and this enables the formation of the circulating ionizing beams.

Our model provides an important link between planetary nebulae and symbiotic nebulae. The fact that these two types of nebulae may be related in their shaping mechanisms (and that the nebulae are sometimes misclassified) has been known for some time (e.g. Morris 1987, 1990; Corradi 1995; Corradi & Schwarz 1995; Schwarz & Corradi 1992; Corradi et al. 2000). Furthermore, our proposed model for M2–9 emphasizes the important role played by binaries in the shaping of nebulae in general, and in the formation of jets in particular. The formation of collimated jets in some of these systems, as evidenced, for example, by the recent observations of He2–90 (Sahai & Nyman 2000), provides important clues for the processes by which astrophysical jets are accelerated and collimated in general (Livio 2000).

Our dynamical model is presented in §2, the properties of the ionizing beams are discussed in §3, and a summary and discussion follow.

2. AN INTERACTING WINDS MODEL

We consider the following configuration for M2–9, which qualitatively describes D-type symbiotic systems. D-type symbiotic systems are wide binaries (orbital periods ~ 100 yrs) in which a white dwarf accretes from the wind of an AGB star or a Mira variable. We propose that in M2–9 a WD companion accretes mass from the wind of an AGB or post-AGB star. An accretion disk is formed, and this disk powers a mildly collimated fast wind (CFW), or a jet (see e.g. Livio 1999, 2000 for a review of jet formation). The general geometry is presented schematically in Figure 1b. For a typical post-AGB star of $0.6 M_{\odot}$ and a WD companion of $0.8 M_{\odot}$, and an orbital period of 120 years (DBCS), the orbital separation is 27 AU. We assume a circular orbit, since an eccentric orbit would have caused the nebula to possess a strong departure from axisymmetry (Soker, Rappaport & Kiger 2001). For a slow wind velocity of 15 km s^{-1} the Bondi-Hoyle accretion rate is ~ 0.01 of the mass loss rate by the post-AGB star. The latter is $\sim 10^{-6} M_{\odot} \text{ yr}^{-1}$ in the case of M2–9 (Lim & Kwok 2000), assuming a spherical, slow wind. The condition for the formation of an accretion disk around the WD can be expressed in the form $R_d \gtrsim R_{\text{WD}}$, where R_d is the outer

radius of the disk that would form given the wind’s specific angular momentum. The disk radius is given approximately by (e.g. Livio & Warner 1984)

$$R_d \simeq 10^{10} \eta^2 \left(\frac{M_{\text{tot}}}{2 M_\odot} \right) \left(\frac{a}{10^{15} \text{ cm}} \right)^{-3} \left(\frac{M_{\text{WD}}}{M_\odot} \right)^3 \left(\frac{V_{\text{rel}}}{20 \text{ km s}^{-1}} \right)^{-8} \text{ cm} , \quad (1)$$

where M_{tot} is the total system mass, a is the separation, V_{rel} is the relative velocity between the WD and the wind, and η is a dimensionless parameter determined by the existing density or velocity gradients in the wind. Typically $\eta \sim 0.25$ (e.g. Ruffert 1999), which for the assumed parameters gives $R_d/R_{\text{WD}} \sim 25$. Thus, for the assumed parameters a disk can form, but the collimation process may not be extremely efficient, since efficient collimation requires $R_d/R_{\text{WD}} \gg 1$ (e.g. Spruit 1996; Livio 1999). If we assume collimation to take place by a poloidal magnetic field [of the form $B_z \sim (r^2/R_{\text{WD}}^2 + 1)^{-1/2}$] threading the disk, then the minimum jet opening angle is given by $\theta \gtrsim (R_{\text{WD}}/R_d)^{1/2} \gtrsim 10^\circ$. This conclusion is quite consistent with the geometry of the inner lobes in M2–9, which gives an opening angle (or collimation angle, defined from the symmetry axis) of the CFW of $\theta_c \simeq 35^\circ$. We assume that roughly 1% of the accreted mass is blown into the CFW (this is consistent with low-efficiency disk outflows; Livio 1999). For the CFW velocity we take $v_f = 5,000 \text{ km s}^{-1}$ (of the order of the Keplerian velocity near the WD surface and consistent with observations; e.g. Livio 2000, Balick 1989). We find from these values, and from the assumption that the mass loss rate from the AGB, or post AGB, star in the equatorial plane is somewhat higher than that along the polar directions (SR00) that the ratio of the slow wind momentum flux (ρv^2) to that of the CFW is ~ 5 , hence the CFW will be bent and the slow wind will flow along and near the symmetry axis. For the assumed parameters, the slow wind impacts the CFW at a large angle, and the bending is less efficient than in the case $a \gg h$ (where h is the height of the impact point where substantial bending occurs; Fig. 1b). Bending to the asymptotic angle θ_d will therefore occur at a height $h = \text{a few} \times h_0 \sim 100 \text{ AU}$ where $h_0 \sim 21 \text{ AU}$ is given by eq. 12 of SR00 (see Fig. 1b).

The CFW passes through a strong shock and forms a hot bubble which compresses and accelerates the slow wind with which it collides, thus forming a dense shell (e.g. SR00, Fig. 1b). It is important to note though that by “shell” we do not refer to a closed elliptical or spherical shell, but rather to a “corkscrew” structure, or a “snail” type structure. Because of the relatively low mass loss rate from the post-AGB star, even the dense shells are rather tenuous, and hence, too faint to be clearly observed. If we assume that all the kinetic energy of the CFW is transferred to the slow wind material along the lobes, which for our assumed collimation angle is of the order of $\epsilon \sim 0.2$ of the slow wind mass, we find the shell velocity to be

$$v_{\text{shell}} \simeq v_s \left[1 + \frac{\dot{M}_f v_f^2}{\epsilon |\dot{M}_s| v_s^2} \right]^{1/2} \simeq 190 \left(\frac{\dot{M}_f}{0.001 \epsilon |\dot{M}_s|} \right)^{1/2} \left(\frac{v_f}{400 v_s} \right) \left(\frac{v_s}{15 \text{ km s}^{-1}} \right) \text{ km s}^{-1}. \quad (2)$$

However, the acceleration of a dense shell by a more dilute hot bubble is prone to Rayleigh-Taylor instabilities, and the shell most likely will break up into many clumps. This has three important effects. First, the acceleration is much less efficient since the hot, shocked CFW leaks between

these clumps. The terminal shell (clump) velocity may thus be as low as $v_{\text{clump}} \sim 50 \text{ km s}^{-1}$. Second, the clumpy shells (a shell each orbital period) allow more of the ionizing radiation to reach the lobes. Finally, the leakage of the CFW keeps the directionality of the expanding bubble to be mainly radial. The clumps reach the bright rings on the lobes (where the N2 and S2 knots are located; see TGD) on a timescale of

$$t_c = 500 \left(\frac{D_r}{0.025 \text{ pc}} \right) \left(\frac{v_{\text{clump}}}{50 \text{ km s}^{-1}} \right)^{-1} \text{ yrs}, \quad (3)$$

where D_r is the distance to the bright rings in the lobes. The dynamical age of M2–9 is $\sim 1,200$ yrs (Schwarz et al. 1997), hence the first shell reached the ring ~ 700 years ago. For an orbital period of 120 years this means that ~ 6 shells have reached the rings. We propose that the high density material in the rings was formed by these clumps (see also TGD).

The following thing should be noted in regard to the mass loss from the WD. As the WD accretes from the wind, it could undergo periodic symbiotic nova outbursts (e.g. Livio, Pringle & Regev 1989). Phases of a strong WD wind could follow such outbursts (e.g. Kovetz 1998 and references therein). In fact, all recorded symbiotic nova eruptions were followed by an intensive Wolf-Rayet type wind from the hot companion (Mikolajewska 2000). The recurrence time for such outbursts is given approximately by (Livio 1994)

$$\tau_{\text{rec}} \simeq 8.7 \times 10^3 \left(\frac{\dot{M}_{\text{acc}}}{10^{-8} \text{ M}_{\odot} \text{ yr}^{-1}} \right)^{-1} \times \left[1.54 \left(\frac{M_{\text{WD}}}{\text{M}_{\odot}} \right)^{-7/3} - 2.0 \left(\frac{M_{\text{WD}}}{\text{M}_{\odot}} \right)^{-1} + 0.65 \left(\frac{M_{\text{WD}}}{\text{M}_{\odot}} \right)^{1/3} \right]^{0.7} \text{ yrs} . \quad (4)$$

Thus, for a $0.8 \text{ M}_{\odot}$ WD we can expect strong WD wind phases roughly every ~ 6700 yrs. We propose that such an outburst formed the “ansae” of M2–9 ~ 1200 years ago. The long time that has elapsed since the eruption implies, in the case of M2–9, a current lower mass loss rate and a faster velocity than in typical post eruption symbiotics. We note that the formation of jets or any ballistic outflows, in short bursts, leads to a linear relation between the distance from the center and the outflow velocity. Such a linear relation is observed in many bipolar PNe, suggesting that bursts are a common phenomenon in bipolar PNe.

Another important aspect of symbiotic-nova outbursts is the fact that following eruptions, the WD enters a supersoft x-ray phase, in which it is characterized by an effective temperature of $150,000\text{--}600,000 \text{ K}$ (e.g. DiStefano & Rappaport 1994; Yungelson et al. 1996). During this phase, the ionization balance of the circumbinary material is determined by the radiation from this embedded supersoft x-ray source (SXS). Fast collimated outflows have been observed in three of the SXSs: RX J0513.9–6951 (3800 km s^{-1} , Southwell et al. 1996; Southwell, Livio, & Pringle 1997); RX J0019.8+2156 ($\sim 1000 \text{ km s}^{-1}$, Becker et al. 1998); and RX J0925.7–4758 (5200 km s^{-1} , Motch 1998). The mass transfer rates in these systems are not known precisely, but are probably above $10^{-8} \text{ M}_{\odot} \text{ yr}^{-1}$ in all three sources.

In principle, the WD can also be a more permanent SXS, similar in nature to systems like CAL 83 and CAL 87. In these systems, the accretion rate onto the WD is sufficiently high for the hydrogen to burn steadily (e.g. van den Heuvel et al. 1992). For a WD of 0.6–0.8 $M_\odot$, the accretion rate for steady burning is of the order of (a few) $\times 10^{-8} M_\odot \text{ yr}^{-1}$ (e.g. Nomoto 1982), definitely achievable by Bondi-Hoyle accretion from the post AGB star’s wind. Steady burning will result in a soft x-ray luminosity of

$$L_x \simeq 10^{37} \left(\frac{\dot{M}_{\text{AGB}}}{10^{-6} M_\odot \text{ yr}^{-1}} \right) \left(\frac{M_{\text{WD}}}{M_\odot} \right)^2 \left(\frac{a}{27 \text{ AU}} \right)^{-2} \left(\frac{V_W}{15 \text{ km s}^{-1}} \right)^{-4} \text{ erg s}^{-1} , \quad (5)$$

and an effective temperature of

$$T_{\text{SXS}} \simeq 5 \times 10^5 \left(\frac{L_x}{10^{37} \text{ ergs s}^{-1}} \right)^{1/4} \text{ K} . \quad (6)$$

3. THE IONIZING “BEAMS”

In order for the soft x-rays to reach the outer visible parts of the M2–9 nebula (and cause them to fluoresce), they must ionize their own path through the stellar wind of the AGB star and other intervening matter, and/or have a path cleared by the action of the CFW coming from the white dwarf. Given that much of the outer nebula apparently receives a substantial ionizing flux, we conclude that the radiation is able to ionize its way through the stellar wind in many if not most directions. A number of works have addressed the question of the ionization structure and concomitant line emission from an ionizing source embedded in a stellar wind (see, e.g. Nussbaumer & Vogel 1987; Seaquist, Taylor & Button 1984). Here we are mainly concerned with the fraction of the radiation flux, as a function of direction, that can pass through the wind and which is available to illuminate the outer nebula. This radiation flux can be estimated as follows.

Even though the ionizing radiation is emitted by the WD companion, while the slow wind is blown by the AGB or post-AGB star, the geometry can still be safely assumed to be nearly spherical. This is a consequence of the fact that the CFW clears the slow wind close to the WD, up to a distance of the order of the bending distance (see previous section), of $r_{\text{min}} \sim 100 \text{ AU}$, which is considerably larger than the orbital separation of $a \sim 27 \text{ AU}$. For the ionizing radiation to be able to reach the lobes, it needs to ionize both the newly ejected slow wind material, and the recombining wind, in its way. For the parameters of the flow discussed here the newly ejected material is not important. The total recombination rate from a minimum distance r_{min} to the lobes at $r_l \gg r_{\text{min}}$ is given by (assuming a spherical flow)

$$\dot{R} = 4\pi \int_{r_{\text{min}}}^{r_l} \alpha n_i n_e r^2 dr = 8 \times 10^{45} \left(\frac{\dot{M}_{\text{AGB}}}{10^{-6} M_\odot \text{ yr}^{-1}} \right)^2 \left(\frac{v_s}{15 \text{ km s}^{-1}} \right)^{-2} \left(\frac{r_{\text{min}}}{100 \text{ AU}} \right)^{-1} \text{ s}^{-1} , \quad (7)$$

where α is the recombination coefficient, n_e and n_i are the electron and ion number densities (respectively). Here we assumed a fully ionized nebula of solar composition, and for the density we used $\rho = \dot{M}_{\text{AGB}}[4\pi r^2 v_s]^{-1}$. The number of ionizing photons emitted by the WD per unit time, assuming a temperature of $T_{\text{SXS}} \simeq 5 \times 10^5$ K, is

$$\dot{S} \simeq 4 \times 10^{46} \left(\frac{L_x}{10^{37} \text{ erg s}^{-1}} \right) \text{ s}^{-1}. \quad (8)$$

The two rates obtained above, for $\dot{R}$ and $\dot{S}$, are within an order of magnitude from each other. This implies that a substantial fraction, but not all, of the radiation will be absorbed along directions facing the AGB star, while only a negligible fraction will be absorbed along directions cleared by the CFW. The CFW clears away the slow wind along its path. If the compressed shell moves at $\sim 50 \text{ km s}^{-1}$ (eq. 2 and following discussion), then during about half an orbital period (60 years), before fresh slow wind material enters the swept-up region, it will move to a distance of $r \sim 600 \text{ AU}$. This means that the total number of recombinations along that direction is much smaller than the number of ionizing photons emitted in that direction. Along other directions, the slow wind’s closest distance to the WD will be less than the distance of bending, and a substantial fraction of the ionizing flux will be absorbed, but some will still reach the lobes. This is consistent with deductions from observations (TGD). The estimates presented by eqs. (7) and (8) therefore support our proposed scenario, in which the east-west motion of the bright knots is caused by ionizing beams which pass through channels cleared by bent “jets” (or a CFW).

4. DISCUSSION AND SUMMARY

M2–9 is a unique nebula in that it shows a variability on a time scale which is consistent with the orbital period expected from the binary model for the formation of bipolar nebulae. In this model most (although not all) bipolar nebulae are formed by binary systems with orbital periods in the range of $\sim 1\text{--}10^3$ yrs (e.g. SR00). The time varying features exhibit a mirror symmetry, rather than a point-symmetry (TGD; DBCS). As we have shown, a mirror symmetry is expected from the interaction of two winds, a slow wind emitted by the AGB or post AGB star, and a jet (or collimated fast wind [CFW]) emanating from the accretion disk around the WD companion. The mirror symmetry is produced as the CFW is pushed and bent by the slow wind. Our model identifies the evolutionary state of M2–9 as one which relates three different classes of binary systems: supersoft X-ray sources (SXSs), symbiotic nebulae, and PNe (or proto-PNe). The ionizing source is the white dwarf companion, and not the mass-losing star (as already concluded by e.g. Goodrich 1991; Schwarz et al. 1997; Solf 2000; Lim & Kwok 2000). At the same time, the fast outward moving “ansae” (Schwarz et al. 1997) and the large and bright nebula strongly suggest that in the recent past the system has both undergone a more intense mass loss episode, and an

outburst associated with the accreting WD companion. Hence, our evolutionary scenario for the nebula can be characterized as a symbiotic nebula evolving to a PN.

While M2–9 is unique in its time variation, it is not in its morphology. The bipolar proto PN Henize 401, for example, has a similar structure (Sahai, Bujarrabal, & Zijlstra 1999), but with no evidence for circulating ionizing beams. We note that Sahai et al. (1999) argue that the bipolar structure in He 401 was formed by collimated bipolar jets, and not by the more commonly assumed interacting winds model, in which a spherical fast wind catches up with a previously ejected bipolar slow wind (e.g. Balick 1987). The model of Sahai et al. has in fact some features that are similar to those of our model for M2–9.

Our model offers an interpretation to the bright knots that is different from the one proposed by DBCS. We showed that the bright knots are exposed to a higher flux of ionizing radiation (from the hot WD) than the rest of the lobes, by virtue of the path cleared by the CFW. DBCS, on the other hand, attribute the brightness/ionization of the knots to shock heating, caused by jets impacting on lobe walls. We see a few problems with this suggestion: (i) it offers no clear explanation for the mirror symmetry (the jets will collide with slow wind material ejected half a period earlier and will be slowed down substantially). (ii) The fact that the off-knots regions are ionized as well (TGD) argues for an ionizing source that is not confined to the small cross section of a jet, which is present at all times. (iii) We find the ‘corkscrew’ model of DBCS (see e.g. their Fig. 2) to provide for a rather poor description of the nebula (e.g. the lines that are supposed to represent the emission in 1989 and 1999 do not follow very well the actual emission). We rather interpret the 1989 and 1999 images as follows. The bright emission regions (from the center of the nebula to the bright blobs along the symmetry axis) are moving in the same sense, with the 1989 emission being toward the outskirts (east side) relative to the 1999 emission. The change in the east-west location occurs only near the bright blobs (N_3 and S_3), and even there it represents a complicated flow structure, and radiative transfer from these regions outward. Nevertheless, our model does share some common features with that of DBCS (e.g. a fast outflow from the companion).

One point in our model deserves further discussion. A cleared path in a certain direction for the ionizing beam (and therefore enhanced ionizing radiation), could result, in principle, in a more extended, rather than brighter, nebula (which is not observed). We should therefore check that for the parameters of M2–9 there is no significant penetration of the ionization front in the direction *perpendicular* to the lobe walls. For an electron density in the lobes of $n_e \sim 10^4 \text{ cm}^{-3}$ (DBCS), an ionizing source of $\dot{S} \sim 4 \times 10^{46} \text{ s}^{-1}$ (eq. 8), and a path clearing for about 10% of the period, one can easily show that the penetration of the ionization perpendicular to the wall would be by less than $\sim 0.1D$ (where D is the radial distance to the S_2 , N_2 knots). Thus, while we do not expect any significant perpendicular penetration, we do expect the bright regions to extend *along* the lobes (due to the lower density), as observed (Fig. 1a).

To conclude, the spectacular nebula M2–9 provides an opportunity to identify an evolutionary link among three types of (empirically defined) binaries: PNe with binary central stars, symbiotic binaries, and supersoft x-ray sources.

ACKNOWLEDGMENTS: ML acknowledges support by NASA Grant NAG5-6857. NS acknowledges partial support by a grant from the Israel Science Foundation, and a grant from the US-Israel Binational Science Foundation. We acknowledge extremely helpful comments and suggestions from Saul Rappaport.

REFERENCES

- Allen, D. A. & Swings, J. P. 1972, ApJ, 174, 583.
- Balick, B. 1987, AJ, 94, 671.
- Balick, B. 1989, AJ, 97, 476.
- Becker, C. M., Remillard, R. A., Rappaport, S., & McClintock, J. E. 1998, ApJ, 506, 880.
- Corradi, R. L. M. 1995, MNRAS, 276, 521.
- Corradi, R. L. M., Livio, M., Schwarz, H. E., & Ulisse, M. 2000, in *Asymmetrical Planetary Nebulae II: From Origins to Microstructures*, eds. J. Kastner, S. Rappaport, and N. Soker, ASP Conference Series Vol. 199, p. 175.
- Corradi, R. L. M. & Schwarz, H. E. 1995, A&A, 293, 871.
- DiStefano, R. & Rappaport, S. 1994, ApJ, 437, 733.
- Doyle, S., Balick, B., Corradi, R. L. M., & Schwarz, H. E. 2000, AJ, 119, 1339 (DBCS).
- Goodrich, R. W. 1991, ApJ, 366, 163.
- Hora, J. L. & Latter, W. B. 1994, ApJ, 437, 281.
- Kohoutek, L., & Surdej, J. 1980, A&A, 85, 161.
- Kovetz, A. 1998, ApJ, 495, 401.
- Lim, J. & Kwok, S. 2000, in *Asymmetrical Planetary Nebulae II: From Origins to Microstructures*, eds. J. Kastner, S. Rappaport, and N. Soker, ASP Conference Series Vol. 199, p. 259.
- Livio, M. 1994, in *Interacting Binaries*, eds. H. Nussbaumer & A. Orr (Berlin: Springer-Verlag), 135.
- Livio, M. 1999, Phys. Rep., 311, 225.
- Livio, M. 2000, in *Cosmic Explosions*, AIP Conf. Proc. 522, eds. S. S. Holt & W. W. Zhang (Melville: AIP), 275.
- Livio, M., Prialnik, D., & Regev, O. 1989, ApJ, 341, 299.
- Livio, M. & Warner, B. 1984, Observatory, 104, 152.
- Mikolajewska, J. 2000, in *Asymmetrical Planetary Nebulae II: From Origins to Microstructures*, eds. J. H. Kastner, N. Soker, and S. Rappaport, ASP Conference Series, vol. 199, p. 431.
- Morris, M. 1987, PASP, 99, 1115.
- Morris, M. 1990, in *From Miras to Planetary Nebulae: Which Path for Stellar Evolution?* eds. M. O. Mennessier & A. Omont (Paris: Editions Frontieres), p. 520.
- Motch, C. 1998, A&A, 338, L13.
- Nomoto, K. 1982, ApJ, 253, 798.

- Nussbaumer, H. & Vogel, M. 1987, A&A, 182, 51.
- Phillips, J. P. & Cuesta, L. 1999, AJ, 118, 2919.
- Ruffert, M. 1999, A&A, 346, 861.
- Sahai, R., Bujarrabal, V., & Zijlstra, A. 1999, ApJ, 518, L115.
- Sahai, R. & Nyman, L.-A. 2000, ApJ, in press.
- Scarrott, R. M. J., Scarrott, S. M., & Wolstencroft, R. D. 1993, MNRAS, 264, 740.
- Schwarz, H. E., Aspin, C., Corradi, R. L. M., & Reipurth, B. 1997, A&A, 319, 267.
- Schwarz, H. E. & Corradi, R. L. M. 1992, A&A, 265, L37.
- Sequist, E. R., Taylor, A. R., & Button, S. 1984, ApJ, 284, 202.
- Spruit, H. C. 1996, in *Evolutionary Processes in Binary Stars*, eds. R. A. M. J. Wijers, M. B. Davies, & C. A. Tout (Cambridge: CUP), 249.
- Soker, N. & Rappaport, S. 2000, ApJ, 538, 241 (SR00).
- Soker, N., Rappaport, S., & Kiger, J. 2001, preprint.
- Solf, J. 2000, A&A, 354, 674.
- Southwell, K. A., Livio, M., Charles, P. A., O’Donoghue, D., & Sutherland, W. J. 1996, ApJ, 470, 1065.
- Southwell, K. A., Livio, M., & Pringle, J. E. 1997, ApJ, 478, L29.
- Torres-Peimbert, S. & Arrieta, A. 1998, RevMex AA (Series de Conferencias) 7, 171.
- Trammell, S. R., Goodrich, R. W., & Dinerstein, H. L. 1995, ApJ, 453, 761 (TGD).
- van den Bergh, S. 1974, A&A, 32, 351.
- van den Heuvel, E. P. J., Bhattacharya, D., Nomoto, K., & Rappaport, S. A. 1992, A&A, 262, 97.
- Yungelson, L. R., Livio, M., Truran, J. W., Tutukov, A. V., & Federova, A. 1996, ApJ, 466, 890.


FIGURE CAPTIONS

Fig. 1a: The general structure of M2–9.

Fig. 1b: A schematic drawing of the flow structure (not to scale). The figure shows the plane perpendicular to the equatorial plane, and momentarily containing the two stars. The 3D structure is that of a corkscrew. Drawn on the figure are: the slow wind from the post-AGB star (stream

lines originating from the post-AGB star); stream lines of the pre-shocked CFW; the shock front of the CFW; The hot bubble region (again, a corkscrew structure in 3D); and the dense shell formed from the slow wind material accelerated by the hot bubble, with its Rayleigh-Taylor instabilities.

M2-9



Interacting Winds Scenario

